

POLYNOMIALS

SECTION A: (1 MARK)

- Form a quadratic polynomial whose zeroes are $\frac{2}{3}$ and $-\frac{1}{3}$. **(CBSE 2008)** $(9x^2 - 3x - 2)$
- If -1 is a zero of the polynomial $f(x) = x^2 - 7x - 8$, then find the other zero. **(CBSE 2012)** (8)
- If α and β are the zeroes of the polynomial $2x^2 + 5x + 1$, then what is the value of $\alpha + \beta + \alpha\beta$? (-2)
- If the sum of the zeroes of the polynomial $P(x) = 3k^2 + (2k + 1)x - k + 5$ is equal to the product of the zeroes, then, find the value of k. (k = -6)
- The graph of the polynomial $f(x) = 2x - 5$ is a straight line. At which point does the graph intersect the x-axis? **(CBSE 2012)** $(\frac{5}{2}, 0)$

SECTION B: (2 MARKS)

- For what value of k, (-4) is a zero of the polynomial $x^2 - x - (2k + 2)$? **(CBSE 2009)** K = 9
- If m and n are the zeroes of the polynomials $3x^2 + 11x - 4$, find the value of $\frac{m}{n} + \frac{n}{m}$. **(CBSE 2012)** $(\frac{-145}{12})$
- If the zeroes of the polynomial $x^2 + px + q$ are double in value to the zeroes of the polynomial $2x^2 - 5x - 3$, find the values of p and q. p = -5, q = -6.
- Form a quadratic polynomial whose one zero is $3 + \sqrt{2}$ and the sum of zeroes is 6. $x^2 - 6x + 7$
- If $ax^2 - 7x + c$ has 14 as the sum of the zeroes and also as product of the zeroes, find the value of a and c. a = $\frac{1}{2}$, c = 7. **(HOTS)**

SECTION C: (3 MARKS)

- Find the zeroes of the following polynomials by factorization method and verify the relations between the zeroes and the coefficients of the polynomial.
 - $2x^2 - (1 + 2\sqrt{2})x + \sqrt{2}$ **(EXEMPLAR)** (i) $1/2, \sqrt{2}$
 - $y^2 + \frac{3}{2}\sqrt{5}y - 5$. (ii) $-2\sqrt{5}, \frac{\sqrt{5}}{2}$
- Find the value of a and b so that $8x^4 + 14x^3 - 2x^2 + ax + b$ is exactly divisible by $4x^2 + 3x - 2$. **(CBSE 2011)** a = -7, b = 2.
- If p and q are the zeroes of the polynomial $6y^2 - 7y + 2$, find a quadratic polynomial whose zeroes are $\frac{1}{p}$ and $\frac{1}{q}$. **(CBSE 2011)** $\frac{1}{2}(2y^2 - 7y + 6)$
- On dividing a polynomial $3x^3 + 4x^2 + 5x - 13$ by a polynomial g(x), the quotient and the remainder were $(3x + 10)$ and $(16x - 43)$ respectively. Find g(x). **(CBSE 2011)** $x^2 - 2x + 3$
- If one zero of a polynomial $3x^2 - 8x + 2k + 1$ is seven times the other, find the value of k. **(CBSE 2011)** K = 2/3.

SECTION D: (4 MARKS)

16. Find the other zeroes of the polynomial $P(x) = 2x^4 + 7x^3 - 19x^2 - 14x + 30$, if two of its zeroes are $\frac{3}{2}$ and -5 . $-\sqrt{2}, \sqrt{2}$
(CBSE 2011)
17. Given $\sqrt{2}$ is a zero of the cubic polynomial $6x^3 + \sqrt{2}x^2 - 10x - 4\sqrt{2}$, find the other two zeroes. $\frac{-\sqrt{2}}{2}, \frac{-2\sqrt{2}}{3}$
18. If the polynomial $x^4 - 6x^3 + 16x^2 - 25x + 10$ is divided by another polynomial $x^2 - 2x + k$, the remainder comes out to be $x + a$, find the values of k and a . $k = 5$
 $a = -5$.
19. If the remainder on division of $x^3 + 2x^2 + kx + 3$ by $(x - 3)$ is 21, find the quotient and the value of k . Hence find the zeroes of the cubic polynomial $x^3 + 2x^2 + kx - 18$. $k = -9$
Quotient = $x^2 + 5x + 6$
Zeroes: 3, -2, -3
(EXEMPLAR)
20. If α and β are the zeroes of the polynomial $p(x) = 2x^2 + 5x + k$ satisfying the relation $\alpha^2 + \beta^2 + \alpha\beta = \frac{21}{4}$, then find the value of k . $k = 2$.