Downloaded from www.studiestoday.com

DEFINITE INTEGRALS

Q.1)	(a) $I = \int_0^2 \frac{1}{4+x-x^2} dx$
Sol.1)	(a) $I = \int_0^2 \frac{1}{4+x-x^2} dx$ $Perfect\ square$ $I = -\int_0^2 \frac{1}{x^2-x-4}$ $I = -\int_0^2 \frac{1}{\left(x-\frac{1}{2}\right)^2-\frac{1}{4}-4} dx$ $I = -\int_0^2 \frac{1}{\left(x-\frac{1}{2}\right)^2-\left(\frac{\sqrt{17}}{2}\right)^2} dx$ $I = \int_0^2 \frac{1}{\left(\frac{\sqrt{17}}{2}\right)^2-\left(x-\frac{1}{2}\right)^2} dx$ $I = \frac{1}{2\times\frac{\sqrt{17}}{2}} \times \left(\log\left\|\frac{\frac{\sqrt{17}}{2}+x-\frac{1}{2}}{\frac{\sqrt{17}}{2}-x+\frac{1}{2}}\right\|\right)_0^2$ $I = \frac{1}{\sqrt{17}}\left(log\left\|\frac{\sqrt{17}+2x-1}{\sqrt{17}-2x+1}\right\|\right)_0^2$ $I = \frac{1}{\sqrt{17}}\left[log\left\|\frac{\sqrt{17}+3}{\sqrt{17}-3}\right\| - log\left\|\frac{\sqrt{17}-1}{\sqrt{17}+1}\right\|\right]$ $I = \frac{1}{\sqrt{17}} log\left\|\frac{\sqrt{17}+3}{\sqrt{17}-3} \times \frac{\sqrt{17}+1}{\sqrt{17}-1}\right\|$ $\ldots\ldots\{log\ A - log\ B = log\ (A/B)\}$ $I = \frac{1}{\sqrt{17}} log\left\|\frac{17+\sqrt{17}+3\sqrt{17}+3}{17-\sqrt{17}-3\sqrt{17}+3}\right\|$ $I = \frac{1}{\sqrt{17}} log\left\|\frac{4\sqrt{17}+20}{20-4\sqrt{17}}\right\|$ $I = \frac{1}{\sqrt{17}} log\left\|\frac{\sqrt{17}+5}{5-\sqrt{17}}\right\|$ $I = \frac{1}{\sqrt{17}} log\left\|\frac{(\sqrt{17}+5)(5+\sqrt{17})}{25-17}\right\|$ $\ldots\ldots(rationalize\ )$ $I = \frac{1}{\sqrt{17}} log\left(\frac{17+25+2\sqrt{17}}{8}\right)$ $I = \frac{1}{\sqrt{17}} log\left(\frac{42+2\sqrt{17}}{8}\right)$ $\therefore\ I = \frac{1}{\sqrt{17}} log\left(\frac{21+\sqrt{17}}{4}\right)$ $Ans\ \ldots.$
Q.2)	$I = \int_1^2 \frac{5(x^2-x-1)}{x^2+3x+2} dx$
Sol.2)	$I = 5\int_1^2 \frac{(x^2-x-1)}{x^2+3x+2} dx$ Here , degree of numerator = degree of denominator Express $q = \frac{R}{D}$ $I = 5\int_1^2 \left(1 - \frac{4x+3}{x^2+3x+2}\right) dx$

Copyright © www.studiestoday.com All rights reserved. No part of this publication may be reproduced, distributed, or transmitted in any form or by any means, including photocopying, recording, or other electronic or mechanical methods, without the prior written permission

Downloaded from www.studiestoday.com

Downloaded from www.studiestoday.com

	$= 5\int_1^2 1\,.dx - 5\int_1^2 \frac{4x+3}{x^2+3x+3}dx$ $= 5(x)_1^2 - 5\int_1^2 \frac{4x+3}{(x+1)(x+2)}dx$ $I = 5 - 5\int_1^2 \frac{4x+3}{(x+1)(x+2)}dx$ Let $\frac{4x+3}{(x+1)(x+2)} = \frac{A}{x+1} + \frac{B}{x+2}$ $\Rightarrow 4x + 3 = A(x + 2) + B(x + 1)$ Comp. the coff. of x and constant $4 = A + B$ $3 = 2A + B$ Solving then equation, we get $A = -1$ & $B = 5$ $\therefore\ I = 5 - 5\int_1^2 \frac{-1}{x+2} + \frac{5}{x+2}dx$ $I = 5 - 5\,[-log\lvert x+1\rvert + 5\,log\lvert x+2\rvert]_1^2$ $= 5 - 5[(-log\,3 + 5\,log\,4) - (log\,2 + 5\,log\,3)]$ $= 5 - 5[-log\,3 + 10\,log\,2 + log\,2 - 5\,log\,3]$ $= 5 - 5[11\,log\,2 - 6\,log\,3]$ $I = 5 - 55\,log\,2 + 30\,log\,3$ *Ans ...*
Q.3)	If $\int_a^b x^3\,dx = 0$ and if $\int_a^b x^2\,dx = \frac{2}{3}$. Find value of a & b.
Sol.3)	Consider $\int_a^b x^3\,dx = 0$ $\Rightarrow \left(\frac{x^4}{4}\right)_a^b = 0$ $\Rightarrow \frac{1}{4}[b^4 - a^4] = 0$ $\Rightarrow a^4 = b^4$ $\Rightarrow a = -b$ Consider $\int_a^b x^2\,dx = \frac{2}{3}$ $\Rightarrow \left(\frac{x^3}{3}\right)_a^b = \frac{2}{3}$ $\Rightarrow \frac{1}{3}(b^3 - a^3) = \frac{2}{3}$ $\Rightarrow b^3 - a^3 = 2$ $\Rightarrow (-a)^3 - a^3 = 2$ $\Rightarrow -a^3 - a^3 = 2$ $\Rightarrow -2a^3 = 2$ $\Rightarrow a^3 = -1$

Copyright © www.studiestoday.com All rights reserved. No part of this publication may be reproduced, distributed, or transmitted in any form or by any means, including photocopying, recording, or other electronic or mechanical methods, without the prior written permission

Downloaded from www.studiestoday.com

Downloaded from www.studiestoday.com

	$\Rightarrow a = -1$ Since $b = -a$ $\therefore b = 1$ $\therefore a = -1 \,\&\, b = 1$ Ans....
Q.4)	$I = \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} cos(2x).\log(sin\,x)\,dx$
Sol.4)	$I = \left[\left(\log(sin\,x).\frac{sin(2x)}{2}\right)_{\frac{\pi}{4}}^{\frac{\pi}{2}} - \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \frac{1}{sin\,x}.cos\,x.\frac{sin(2x)}{2}dx\right]$ $= \left[\left(\log 1.\frac{sin\,\pi}{2}\right) - \left(\log\left(\frac{1}{\sqrt{2}}\right).\frac{sin\frac{\pi}{2}}{2}\right)\right] - \frac{1}{2}\int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \frac{1}{sin\,x}.cos\,x.2sin\,x\,cos\,x\,dx$ $= \left[0 + log(\sqrt{2})\frac{1}{2}\right] - \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} cos^2x\,dx$ ……..$\{log\,(a/b) = -log\,(b/a)\}$ $= \frac{1}{2}log\,2\frac{1}{2} - \frac{1}{2}\int_{\frac{\pi}{4}}^{\frac{\pi}{2}} 1 + cos(2x)\,dx$ $= \frac{1}{2}log\,2\,\frac{1}{2} - \frac{1}{2}\left[x + \frac{sin(2x)}{2}\right]_{\frac{\pi}{4}}^{\frac{\pi}{2}}$ $= \frac{1}{2}log\,2\,\frac{1}{2} - \frac{1}{2}\left[\left(\frac{\pi}{2} + 0\right) - \left(\frac{\pi}{4} + \frac{1}{2}\right)\right]$ $= \frac{1}{2}log\,2\,\frac{1}{2} - \frac{\pi}{4} + \frac{\pi}{8} + \frac{1}{4}$ $I = \frac{1}{4}\,log\,2 - \frac{\pi}{8} + \frac{1}{4}$ ans.
Q.5)	$I = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{sin\,x + cos\,x}{\sqrt{sin(2x)}}dx$
Sol.5)	$I = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{sin\,x + cos\,x}{\sqrt{1-1+sin(2x)}}dx$ $= \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{sin\,x + cos\,x}{\sqrt{1-(1-sin\,2x)}}dx$ $= \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{sin\,x + cos\,x}{\sqrt{1-[sin^2x + cos^2x - 2\,sin\,x\,cos\,x]}}dx$ $= \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{sin\,x + cos\,x}{\sqrt{1-(sin\,x - cos\,x)^2}}dx$ $Put\ sin\,x - cos\,x = t$ $when\ x = \frac{\pi}{6}$ $t = \frac{1}{2} - \frac{\sqrt{3}}{2} = \frac{1-\sqrt{3}}{2}$ $(cos\,x + sin\,x)dx = dt$ $when\ x = \frac{\pi}{3}$ $t = \frac{\sqrt{3}}{2} - \frac{1}{2} = \frac{\sqrt{3}-1}{2}$ $\therefore I = \int_{\frac{1-\sqrt{3}}{2}}^{\frac{\sqrt{3}-1}{2}} \frac{dt}{\sqrt{1-t^2}}$ $= (sin^{-1}t)_{\frac{1-\sqrt{3}}{2}}^{\frac{\sqrt{3}-1}{2}}$ $= sin^{-1}\left(\frac{\sqrt{3}-1}{2}\right) - sin^{-1}\left(\frac{1-\sqrt{3}}{2}\right)$

Copyright © www.studiestoday.com All rights reserved. No part of this publication may be reproduced, distributed, or transmitted in any form or by any means, including photocopying, recording, or other electronic or mechanical methods, without the prior written permission

Downloaded from www.studiestoday.com

Downloaded from www.studiestoday.com

	$= sin^{-1}\left(\frac{\sqrt{3}-1}{2}\right) + sin^{-1}\left(\frac{\sqrt{3}-1}{2}\right)$ ……{∵ $sin-1(-x) = -sin-1\,x$} $I = 2\,sin^{-1}\left(\frac{\sqrt{3}-1}{2}\right)$ *Ans* ….
Q.6)	$I = \int_0^{\frac{\pi}{4}} \frac{sin\,x+cos\,x}{9+16\,sin\,(2x)}dx$
Sol.6)	$I = \int_0^{\frac{\pi}{4}} \frac{sin\,x+cos\,x}{9+16\,[1-1+sin(2x)]}dx$ $I = \int_0^{\frac{\pi}{4}} \frac{sin\,x+cos\,x}{9+16\,[1-(1-sin\,2x)]}dx$ $I = \int_0^{\frac{\pi}{4}} \frac{sin\,x+cos\,x}{9+16\,[1-(sin^2\,x+cos^2\,x-2\,sin\,x\,cos\,x)]}dx$ $I = \int_0^{\frac{\pi}{4}} \frac{sin\,x+cos\,x}{9+16\,[1-(sin\,x-cos\,x)^2]}dx$ *Put* $sin\,x - cos\,x = t$ *when* $x = 0, t = 0 - 1 = -1$ $(cos\,x + sin\,x)dx = dt$ *when* $x = \frac{\pi}{4}, t = \frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}} = 0$ $\therefore\ I = \int_{-1}^{0} \frac{dt}{9+16(1-t^2)}$ $= \int_{-1}^{0} \frac{1}{25-16t^2}dt$ $= \frac{1}{16}\int_{-1}^{0} \frac{1}{\left(\frac{5}{4}\right)^2-t^2}dt$ $= \frac{1}{16} \times \frac{1}{2\times\frac{5}{4}}\left(log\left\|\frac{\frac{5}{4}+t}{\frac{5}{4}-t}\right\|\right)_{-1}^{0}$ $= \frac{1}{40}\left(log\left\|\frac{5+4t}{5-4t}\right\|\right)_{-1}^{0}$ $= \frac{1}{40}\left[log\|1\| - log\left\|\frac{1}{9}\right\|\right]$ $= \frac{1}{40}[0 + log(9)]$ $I = \frac{1}{40}log\,9$ $I = \frac{1}{20}log\,3$ *Ans* ….
Q.7)	(a) $I = \int_1^2 e^{2x}\left(\frac{1}{x} - \frac{1}{2x^2}\right)dx$
Sol.7)	(a) $I = \int_1^2 e^{2x}.\frac{1}{x}dx - \frac{1}{2}\int_1^2 e^{2x}.\frac{1}{x^2}dx$ $= \left(\frac{1}{x}.\frac{e^{2x}}{2}\right)_1^2 + \frac{1}{2}\int_1^2 e^{2x}.\frac{1}{x^2}dx - \frac{1}{2}\int_1^2 e^{2x}.\frac{1}{x^2}dx$ $= \left(\frac{1}{2}.\frac{e^4}{2}\right) - \left(\frac{1}{1}.\frac{e^2}{2}\right)$ $= \frac{e^4}{4} - \frac{e^2}{2}$ $= \frac{e^2}{2}\left(\frac{e^2}{2} - 1\right)$ Ans…..

Copyright © www.studiestoday.com All rights reserved. No part of this publication may be reproduced, distributed, or transmitted in any form or by any means, including photocopying, recording, or other electronic or mechanical methods, without the prior written permission

Downloaded from www.studiestoday.com

Downloaded from www.studiestoday.com

Q.8)	$I = \int_0^{\frac{\pi}{2}} \frac{\cos^2 x}{\cos^2 x + 4\sin^2 x} dx$
Sol.8)	Divide N & D by $\cos^4 x$ $I = \int_0^{\frac{\pi}{2}} \frac{\sec^2 x}{\sec^2 x + 4\tan^2 x \sec^2 x} dx$ $I = \int_0^{\frac{\pi}{2}} \frac{\sec^2 x}{\sec^2 x(1+4\tan^2 x)} dx$ $I = \int_0^{\frac{\pi}{2}} \frac{\sec^2 x}{(1+\tan^2 x)(1+4\tan^2 x)} dx$ $\quad$ $put\ \tan x = t$, $\sec^2 x\, dx = dt$, $when\ x = 0\ ;\ t = 0$, $when\ x = \frac{\pi}{2}\ ;\ t = \infty$ $I = \int_0^{\infty} \frac{dt}{(1+t^2)(1+4t^2)} dx$ Type: partial fraction type 4 Let $t^2 = y$ (temp.) $\therefore \frac{1}{(1+t^2)(1+4t^2)} = \frac{1}{(1+y)(1+4y)}$ let $\frac{1}{(1+y)(1+4y)} = \frac{A}{1+y} + \frac{B}{1+4y}$ $1 = A(1 + 4y) + B(1 + y)$ Comp. $0 = 4A + B$ $1 = A + B$ Solving these equations $A = -\frac{1}{3}$ & $B = \frac{4}{3}$ $\therefore I = \int_0^{\infty} \frac{-1}{3(1+t^2)} + \frac{4}{3} \cdot \frac{1}{(1+4t^2)} dt$ $= \frac{-1}{3}\int_0^{\infty} \frac{1}{1+t^2} dt + \frac{4}{3} \cdot \frac{1}{4}\int_0^{\infty} \frac{1}{\frac{1}{4}+t^2} dt$ $= \frac{-1}{3}\int_0^{\infty} \frac{1}{1+t^2} dt + \frac{1}{3}\int_0^{\infty} \frac{1}{\left(\frac{1}{2}\right)^2+t^2} dt$ $= \frac{-1}{3}\tan^{-1}(t)_0^{\infty} + \frac{1}{3} \times 2\left(\tan^{-1}(2t)\right)_0^{\infty}$ $= \frac{-1}{3}(\tan^{-1}\infty - \tan^{-1}0) + \frac{2}{3}[\tan^{-1}(\infty) - \tan^{-1}(0)]$ $= \frac{-1}{3}\left[\frac{\pi}{2}\right] + \frac{2}{3}\left[\frac{\pi}{2} - 0\right]$ $= \frac{-\pi}{6} + \frac{\pi}{3}$ $I = \frac{\pi}{6}$ Ans.....
Q.9)	$\int_0^1 \sqrt{\frac{1-x}{1+x}} dx$
Sol.9)	$\int_0^1 \sqrt{\frac{1-x}{1+x}} dx$

Copyright © www.studiestoday.com All rights reserved. No part of this publication may be reproduced, distributed, or transmitted in any form or by any means, including photocopying, recording, or other electronic or mechanical methods, without the prior written permission

Downloaded from www.studiestoday.com

Downloaded from www.studiestoday.com

	rationalize $I = \int_0^1 \frac{1-x}{\sqrt{1-x^2}} dx$ Separate $I = \int_0^1 \frac{1}{\sqrt{1-x^2}} dx - \int_0^1 \frac{x}{\sqrt{1-x^2}} dx$ $Put\ 1 - x^2 = t \qquad when\ x = 0\ ;\ t = 1$ $x\,dx = -\frac{dt}{2} \qquad when\ x = 1\ ;\ t = 0$ $\therefore I = \int_0^1 \frac{1}{\sqrt{1-x^2}} dx + \frac{1}{2}\int_1^0 \frac{dt}{\sqrt{t}}$ $I = (sin^{-1}x)_0^1 + \frac{1}{2}\left(2\sqrt{t}\right)_1^0$ $I = \left(\frac{\pi}{2} - 0\right) + \frac{1}{2}(0-2)$ $I = \frac{\pi}{2} - 1$ ans.
Q.10)	$I = \int_0^{\frac{1}{\sqrt{2}}} \frac{sin^{-1}x}{(1-x^2)^{\frac{3}{2}}}$
Sol.10)	$I = \int_0^{\frac{1}{\sqrt{2}}} \frac{sin^{-1}x}{(1-x^2)^{\frac{3}{2}}}$ $I = \int_0^{\frac{1}{\sqrt{2}}} \frac{sin^{-1}x}{(1-x^2)\sqrt{1-x^2}} dx$ Put $sin^{-1}x = t \qquad$ when $x = 0\ ;\ t = 0$ $\frac{1}{\sqrt{1-x^2}} dx = dt \qquad$ when $x = \frac{1}{\sqrt{2}}\ ;\ t = \frac{\pi}{4}$ $I = \int_0^{\frac{\pi}{4}} \frac{t}{(1-x^2)} dt$ $= \int_0^{\frac{\pi}{4}} \frac{t}{(1-sin^2 t)} dt \qquad \ldots\ldots\{\because\ x = sin\ t\}$ $= \int_0^{\frac{\pi}{4}} sec^2\ t\,.\,t\ dt$ $= (t\ tan\ )_0^{\frac{\pi}{4}} - \int_0^{\frac{\pi}{4}} 1\,.\,tan\ t\ dt$ $= \left(\frac{\pi}{4}.\,tan\frac{\pi}{4} - 0\right) - (\log\lvert sec\ t\ \rvert)_0^{\frac{\pi}{4}}$ $= \frac{\pi}{4} - \left[log(\sqrt{2}) - log(1)\right] \qquad \ldots\ldots\left\{\because\ sec\frac{\pi}{4} = \sqrt{2} sec\ \theta = 1\right\}$ $= \frac{\pi}{4} - \left[\frac{1}{2}\log 2\right]$ $I = \frac{\pi}{4} - \frac{1}{2}\log 2$ Ans...

Copyright © www.studiestoday.com All rights reserved. No part of this publication may be reproduced, distributed, or transmitted in any form or by any means, including photocopying, recording, or other electronic or mechanical methods, without the prior written permission

Downloaded from www.studiestoday.com