


Class 12th
Relations & Functions

Q.1) Let A and B are any two-empty sets. Show that $f: A \times B \rightarrow B \times A$ such that $f(a, b) = (b, a)$ is a bijective function.

Sol.1) (Rough diagram)



One-One function

let (a, b) and $(c, d) \in A \times B$ (domain)

and $f(a, b) = f(c, d)$

$$\Rightarrow (b, a) = (d, c)$$

$$\Rightarrow b = d \text{ and } a = c$$

$$\Rightarrow (a, b) = (c, d)$$

$\therefore f$ is one-one function

On-To

(.) since $n(A \times B) = n(B \times A)$

(.) f is one-one (just proved above)

(.) \square Range = co-domain

$\therefore f$ must be on o

$\therefore f$ is a bijective function

ans.


Q.2) Show that $f: N \rightarrow N$ given by

$$f(x) = \{x + 1 ; \text{if } x \text{ is odd}\}$$

$$= \{x - 1 ; \text{if } x \text{ is even}\}$$

f is a bijective function.

Sol.2) (Rough diagram)



One-One function :-

(.) Case 1: let x_1 , and x_2 on both odd

$$x_1, x_2 \in N \text{ (domain)}$$

$$\text{and } f(x_1) = f(x_2)$$

$$\Rightarrow x_1 + 1 = x_2 + 1$$

$$\Rightarrow x_1 = x_2$$

(.) Case 2: let x_1 , and x_2 both even

$$x_1, x_2 \in N \text{ (domain)}$$



and $f(x_1) = f(x_2)$
 $\Rightarrow x_1 - 1 = x_2 - 1$
 $\Rightarrow x_1 = x_2$
 (.) Case 3 : let x_1 is odd and x_2 is even
 and $f(x_1) = f(x_2)$
 $\Rightarrow x_1 + 1 = x_2 - 1$
 $\Rightarrow x_2 - x_1 = 2$ {not possible \because even no – odd no $\neq 2$ }
 \therefore thus case is rejected
 (.) Case 4 : let x_1 is even and x_2 is odd
 $f(x_1) = f(x_2)$
 $\Rightarrow x_1 - 1 = x_2 + 1$
 $\Rightarrow x_1 - x_2 = 2$ (not possible \because even no – odd no $\neq 2$)
 \therefore thus case is also rejected

Hence, overall f is one-one function

On-To :

For every odd number $(2n - 1) \in N$ (co-domain) there exists an even number $(2n)$ in domain (N)
 and for every even number $(2p) \in N$ (co-domain) there exists an odd number

$(2p - 1) \in N$ (domain)

\Rightarrow co-domain = Range

$\therefore f$ is on-to

$\therefore f$ is bijective function

Q.2) Given examples of two functions $f : N \rightarrow Z$ and $g : Z \rightarrow Z$ such that gof is injective but g is not injective.

Sol.2) Given : $f : N \rightarrow Z$

and $g : Z \rightarrow Z$

then domain of ' gof ' is same as domain of ' f ' and co-domain of ' gof ' is as same as co-domain of ' g '

$\therefore \text{gof} : N \rightarrow Z$

let $f(x) = x$ and $g(x) = |x|$

$\text{gof} = g(f(x))$

$= g(x)$

$\text{gof} = |x|$

one-one (for gof)

let $x_1, x_2 \in N$ (domain of gof)

and $(\text{gof})(x_1) = (\text{gof})(x_2)$

$\Rightarrow g(f(x_1)) = g(f(x_2))$

$\Rightarrow |x_1| = |x_2|$

$\Rightarrow x_1 = \pm x_2$

but $x_1 \neq x_2$ { $\because x_1, x_2 \in N$ }

$\therefore x_1 = x_2$

$\therefore \text{gof}$ is one-one function

Now $g(-1) = |-1| = 1$

$g(1) = |-1| = 1$

since two different elements in domain (z) of g has same image in co-domain (z)

$\therefore g$ is not one-one



$$\therefore f(x) = x \text{ and } g(x) = |x| \text{ ans.}$$

Q.3) (i) $f(x) = (3 - x^3)^{\frac{1}{3}}$. Find $f \circ f(x)$

Sol.3) (i) $f \circ f = f(f(x))$
 $= f[(3 - x^3)^{\frac{1}{3}}]$
 $= [3 - ((3 - x^3)^{\frac{1}{3}})^3]^{\frac{1}{3}}$
 $= [3 - (3 - x^3)]^{\frac{1}{3}}$
 $= [3 - 3 + x^3]^{\frac{1}{3}}$
 $= (x^3)^{\frac{1}{3}}$
 $= x$
 $\therefore f \circ f = x \text{ ans.}$

(ii) $f(x) = |x|$

$g(x) = |5x - 2|$

Is $f \circ g = g \circ f$ for all $x \in \mathbb{R}$?

$f \circ g = f(g(x))$
 $= f(|5x - 2|)$
 $= ||5x - 2||$
 $= |5x - 2| \quad \{\dots ||x|| = |x|\}$

$g \circ f = g(f(x))$
 $= g(|x|)$
 $= |5|x| - 2|$

clearly $f \circ g \neq g \circ f$ ans.

e.g when $x = -1$

$f \circ g = |5(-1) - 2| = |-5 - 2| = |-7| = 7$

$g \circ f = |5|-1|-2| = |5 - 2| = 3$

(iii) If $f(x) = 2x$; $g(y) = 3y + 4$ and $h(z) = \sin z$

Show that $h \circ (g \circ f) = (h \circ g) \circ f$

LHS = $h \circ (g \circ f)$
 $= h \circ [g(f(x))]$
 $= h \circ [g(2x)]$
 $= h \circ [3(2x) + 4]$
 $= h \circ (6x + 4)$
 $= \sin(6x + 4)$

RHS = $(h \circ g) \circ f$
 $= [h \circ g] \circ f$
 $= [h(g(y))] \circ f$
 $= [(3y + 4)] \circ f$
 $= [\sin(3y + 4)] \circ f$
 $= \sin(3y + 4) \circ (2x)$
 $= \sin(3(2x) + 4) = \sin(6x + 4)$

$\therefore \text{LHS} = \text{RHS}$

Q.4) Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be defined as $f(x) = 10x + 7$. Find function $g(x)$ such that $f \circ g = g \circ f = I_{\mathbb{R}}$

Copyright © www.studiestoday.com All rights reserved. No part of this publication may be reproduced, distributed, or transmitted in any form or by any means, including photocopying, recording, or other electronic or mechanical methods, without the prior written permission



(where $I_R = x$ identity function : $R \rightarrow$ real no's)

Sol.4) We have $f(x) = 10x + 7$
 given $f \circ g = g \circ f = I_R$
 $\Rightarrow f \circ g = g \circ f = x$ where $x \in R$

To find : $g(x)$:

$$\begin{aligned}\text{Consider } f \circ g &= x \\ \Rightarrow f(g(x)) &= x \\ \Rightarrow 10g(x) + 7 &= x \\ \Rightarrow g(x) &= \frac{x-7}{10}\end{aligned}$$

$$\begin{aligned}\text{Now } g \circ f &= g(f(x)) \\ &= g(10x + 7) \\ &= \frac{10x + 7 - 7}{10} \\ &= \frac{10x}{10} = x = I_R \quad (\text{verified}) \\ \dots g(x) &= \frac{x-7}{10} \quad \text{ans.}\end{aligned}$$

Q.5) Let $f : R \rightarrow R$ be the sign um function defined as

$$\begin{aligned}f(x) &= \{-1 ; x < 0\} \\ &\{0 ; x = 0\} \\ &\{1 ; x > 0\}\end{aligned}$$

and $g(x) = [x]$ be the greatest integer function. Then does $f \circ g$ and $g \circ f$ coincide (equal) in $(0, 1)$?

Sol.5) When $x \in (0, 1)$

value of $g(x) = [x]$ can be 0 or 1 $\{[0.1]=0\} \{[0.2]=0\} \{[1]=1\}$

value of $f(x) = 1$ $\{ \because \text{when } x > 0 f(x) = 1 \}$

$$\begin{aligned}\text{Now } f \circ g &= f(g(x)) \\ &= f([x]) \\ &= f(0 \text{ or } 1) \quad \dots \{ \text{as } x \in [0, 1] [x] \text{ can be } 0 \text{ or } 1 \} \\ &= 0, 1 \quad \dots \{ \text{when } x = 0 ; f(x) = 0, \text{ when } x = 1 ; f(x) = 1 \}\end{aligned}$$

$$\begin{aligned}\text{Now } g \circ f &= g(f(x)) \\ &= g(1) \quad \dots \{ \dots x \in [0, 1], f(x) = 1 \} \\ &= [1] = 1\end{aligned}$$

clearly $f \circ g$ does not coincide (equal) with $g \circ f$ when $x \in [0, 1]$ ans.

Q.6) Let $f : \{1, 3, 4\} \rightarrow \{1, 2, 5\}$ and $g : \{1, 2, 5\} \rightarrow \{1, 3\}$ be given by $f = \{(1, 2), (3, 5), (4, 1)\}$ and $g = \{(1, 3), (2, 3), (5, 1)\}$ write down $g \circ f$.

Sol.6) Domain of $g \circ f$ is same as domain of f and co-domain of $g \circ f$ is same as co-domain of g

$$\therefore g \circ f : \{1, 3, 4\} \rightarrow \{1, 3\}$$

$$\begin{aligned}\text{Now, given : } f(1) &= 2 & g(1) &= 3 \\ f(3) &= 5 & g(2) &= 3 \\ f(4) &= 1 & g(5) &= 1\end{aligned}$$

$$g \circ f(1) = g(f(1)) = g(2) = 3$$

$$g \circ f(3) = g(f(3)) = g(5) = 1$$


$$g \circ f(4) = g(f(4)) = g(1) = 3$$

$$\therefore g \circ f = \{(1, 3), (3, 1), (4, 3)\} \quad \text{ans.}$$



Q.7) Let $A = \{1, 2, 3\}$ and $B = \{4, 5, 6, 7\}$ and $f = \{(1, 4), (2, 5), (3, 6)\}$ be a function from A to B. State whether f is one-one or on-to.

Sol.7) Given $f(1) = 4$ $f(2) = 5$ $f(3) = 6$



Clearly f is one-one, as every element in domain (A) has a unique image in co-domain (B)

Since $7 \in$ co-domain (B), but this is not the image of any element in domain (A)

\therefore f is not on-to ans.

Q.8) Consider the function $f: \left[0, \frac{\pi}{2}\right] \rightarrow R$ given by $f(x) = \sin x$ and $g: \left[0, \frac{\pi}{2}\right] \rightarrow R$ given by $g(x) = \cos x$. Show that f and g are one-one but $f + g$ is not one-one.

Sol.8) We know that for any two different elements x_1 and $x_2 \in \left[0, \frac{\pi}{2}\right]$

$$\sin x_1 \neq \sin x_2 \text{ and } \cos x_1 \neq \cos x_2$$

$$\Rightarrow f(x_1) \neq f(x_2) \text{ and } g(x_1) \neq g(x_2)$$

for all $x_1, x_2 \in \left[0, \frac{\pi}{2}\right]$ and $x_1 \neq x_2$

\therefore f and g are one-one

$$\text{Now } f + g = \sin x + \cos x$$

$$(f + g)(0) = f(0) + g(0) = \sin(0) + \cos(0) = 0 + 1 = 1$$

$$(f + g)\left(\frac{\pi}{2}\right) = f\left(\frac{\pi}{2}\right) + g\left(\frac{\pi}{2}\right) = \sin\left(\frac{\pi}{2}\right) + \cos\frac{\pi}{2} = 1 + 0 = 1$$

$$\text{clearly } (f + g)(0) = (f + g)\left(\frac{\pi}{2}\right)$$

$$\text{but } 0 \neq \frac{\pi}{2}$$

i.e two different elements in domain $\left[0, \frac{\pi}{2}\right]$ has same image in co-domain (R)

\therefore $f + g$ is not one-one ans.

Q.9) (i) If $A = \{1, 2, 3\}$ and $B = \{a, c, d, e\}$. Find number of one-one functions



(ii) Find the number of on-to function from A to A if $A = \{1, 2, 3, \dots, n\}$

Sol.9) (i) The element 1 in A can be attached / associated with any element of B in 4 ways
element 2 in A can be attached / Associated in 3 ways
and element 3 can be associated in 2 ways

$$\therefore \text{ total no. of one-one function} = 4 \times 3 \times 2 = 24 \quad \text{ans.}$$

(ii) The element 1 in co-domain can be attached / Associated with any element of domain in = n



ways


element 2 can be associated in = $(n - 1)$ ways

element 3 can be associated in = $(n - 2)$ ways

element n can be associated in = 1 way

\therefore the total no of on-to function an = $n \times (n - 1) \times (n - 2) \times \dots \times 1 = n!$ ans.

Q.10)



Which of the following graphs represent a function ?

Sol.10) (a) is a function

\therefore for each value of x , $f(x)$ attains a unique and different value.

(b) is not a function

since for same value of x , $f(x)$ has multiple values.